## A Highly Collimated Molecular Hydrogen Jet in IRAS 05487-I 0255.

## NIR Imaging and Spectroscopy

- A. Noriega-Crespo (IPAC), Peter M. Garnavich (CfA), Alejandro C. Raga (IAUNAM)

& Karl-Heinz Böhm (UW)

New narrow band near infrared images at the H<sub>2</sub>-1- () 2.121 µm and continuum bands, toget her with K band spectra of a highly collimated bipolar jet close to the IRAS 05487-10255 source are p resented. The jet is located at ~ 30" 'West of the 1 lerbig-Haro 110 outflow. The jet and counter jet are not visible at optical wavelengths, and therefore, do not fall into the 'standard' Herbig-Haro objects classification scheme. Nevertheless, it belongs to an ever growing group of nolecular hydrogen jets associated with YSOs which are optically undetected. The jet looks very well-collimated, with a length-to-width ratio ~ 10 - 20, and to understand this morphology and emission may require to invoke a process like entrainment in a turbulent mixing layer.

The spectra of the jet and counter-jet in the K-band show a few strong  $\rm H_2$  lines, despite their limited signal-to-noise and the low-surface-brightness of the jet. Based on the 4-strongest lines excitation temperatures of 1604±805 K and 2314±365 14, for the red and blue-shifted jet components respectively, were obtained. The estimated reddening in the blue-shifted jet using the (1,0)-Q(2) to (1,0)-S() line ratio, gives a visual extinction  $\rm A_{V} \sim 14$ ". The lines in the red-shifted component arc-too faint for a reliable determination of its reddening. The radial velocities of the jet and counterjet, based on the shift of the (1,0)-S(1)-2.121 $\mu$ m line, are  $\sim 180$  km s<sup>-1</sup> and  $\sim -275$  km s<sup>-1</sup> respectively, suggesting a relatively large angle ( $\sim 30^{\circ} - 45^{\circ}$ ) between the jet and the line of sight, The H<sub>2</sub> emission of the entire jet extends for at least 32 ' or  $\sim 0.1 pc$  at the distance of Orion. If the flow velocity is comparable to that of the radial velocities, then the dynamical age of the system is quite short ( $\sim 500$  yrs), consistent with a Young jet arising from an embedded source, as is the case e.g. for HH211.